



President

Ehab Y. Hanna, MD
Houston, TX

President Elect

Cherie-Ann Nathan, MD
Shreveport, LA

Vice-President

Bert O'Malley, MD
Philadelphia, PA

Secretary

Brian B. Burkey, MD, MEd
Cleveland, OH

Treasurer

Bevan Yueh, MD
Minneapolis, MN

Past Presidents

Jonathan Irish, MD, MSc,
FRCSC
Toronto, ON

Jeffrey Myers, MD, PhD
Houston, TX

Dennis Kraus, MD, FACS
New York, NY

Division Chairs

Administration Division
William Lydiatt, MD
Omaha, NE

Education Division
David Goldenberg, MD, FACS
Hershey, PA

Patient Care Division
Daniel Deschler, MD, FACS
Boston, MA

Research Division
James Rocco, MD, PhD
Columbus, OH

Foundation Chair

Dennis Kraus, MD, FACS
New York, NY

Administrative Office
11300 W. Olympic Blvd.,
Suite 600
Los Angeles, CA 90064
Phone: (310) 437-0559
Fax: (310) 437-0585
Email: admin@ahns.info
www.ahns.info

Position Statement on Human Papillomavirus (HPV) Vaccination for Prevention of HPV-Related Oropharyngeal Cancer

Endorsed by AHNS December 18, 2018

The rate of new diagnoses of Human Papillomavirus (HPV)-related oropharyngeal (i.e. tonsil and base of tongue) squamous cell carcinoma (OPSCC) has been increasing at a dramatic rate in the United States, with HPV now being observed in over 70% of these tumors^{1,2,3}. Recent data suggests that the incidence of HPV-related OPSCC now exceeds that of HPV-related cervical cancer in the United States⁴. Despite these statistics, public awareness of HPV-related OPSCC remains low⁵.

Since 2006, the Food and Drug Administration (FDA) has approved three commercially available vaccines to prevent HPV infection: Gardasil-4 (Merck & Co.), Cervarix (GlaxoSmithKline), and recently, Gardasil-9 (Merck & Co.). Gardasil-9 is currently recommended by the Centers for Disease Control (CDC) Advisory Committee on Immunization Practices (ACIP) for use in females between 9 to 26 years of age and males between 9 to 21 years of age, with the option to vaccinate up to age 26 for certain high-risk individuals⁶. Moreover, the FDA recently expanded the indications for use in both men and women up to 45 years of age.

These HPV vaccines are safe, with numerous large studies and over 10 years of follow-up demonstrating that severe adverse events are very rare and occur at a rate similar to other adolescent vaccines^{7,8,9,10,11}. In addition, vaccine efficacy for preventing oral HPV infections has been shown to be up to 93%, suggesting the potential to prevent HPV-related OPSCC^{12,13}. While these vaccines have clearly demonstrated the ability to reduce anogenital HPV infection and premalignant lesions, a trial evaluating for prevention of HPV-related OPSCC or even a precursor lesion has not been done. This is primarily due to the inability to reliably screen for oropharyngeal premalignancy, the relative rarity of OPSCC, and the resulting need for prohibitively large numbers of individuals to enroll in such a trial. However, a recent meeting by the International Agency for Research on Cancer and the United States National Cancer Institute suggested that prevention of persistent oral infection of 6 months or longer may be an appropriate endpoint to consider for vaccine indication¹⁴.

Despite the safety of these vaccines and their potential benefit, uptake in the United States, especially in males, has been poor^{13,15,16,17}, with lack of physician recommendation and low public awareness being contributing

factors^{18,19}. Low male vaccination rates are particularly relevant since males make up the vast majority of patients who develop HPV-related OPSCC^{1,20}.

Without a definitive change to current HPV immunization practices, the consensus is that the recent trends in HPV-related OPSCC will continue. Therefore, based on the observed link between HPV infection and OPSCC and the safety and efficacy shown of the currently available HPV vaccines in preventing oral HPV infection, The American Head and Neck Society, The American Academy of Otolaryngology – Head and Neck Surgery and The Head and Neck Cancer Alliance strongly encourage HPV vaccination of both boys and girls for prevention of OPSCC and anogenital cancers. We hope that this endorsement will result in more widespread implementation of these vaccines, especially in males, and thus decrease the future burden of HPV infection and HPV-related OPSCC.

Acknowledgements: This document was prepared by the Cancer Prevention Service of the AHNS. The AHNS would also like to recognize Drs. Cherie-Ann Nathan, Vikas Mehta, Terry Day, Carole Fakhry, and Wendell Yarbrough for their contributions to this work.



Daniel G. Deschler, MD, FACS
Chair, Patient Care Division
The American Head and Neck Society



Derrick T. Lin, MD, FACS
Chair, Head and Neck Surgery Committee
The American Academy of Otolaryngology-Head and Neck Surgery



Michael G. Moore, MD, FACS
President, The Head and Neck Cancer Alliance
Chair, Cancer Prevention Service of the American Head and Neck Society

¹ Chaturvedi, A. K. et al. Human papillomavirus and rising oropharyngeal cancer incidence in the United States. *J Clin Oncol* **29**, 4294-4301 (2011).

² Sturgis, E. M. & Cinciripini, P. M. Trends in head and neck cancer incidence in relation to smoking prevalence: an emerging epidemic of human papillomavirus-associated cancers? *Cancer* **110**, 1429-1435 (2007).

³ Saraiya, M. et al. US Assessment of HPV Types in Cancers: Implications for Current and 9-Valent HPV Vaccines. *J Natl Cancer Inst* **107**, (2015).

⁴ Siegel, R. L., Miller, K. D. & Jemal, A. Cancer statistics, 2015. *CA Cancer J Clin* **65**, 5-29 (2015).

-
- ⁵ Luryi, A. L. et al. Public awareness of head and neck cancers: a cross-sectional survey. *JAMA Otolaryngol Head Neck Surg* **140**, 639-646 (2014).
- ⁶ Meites E, Kempe A, Markowitz LE. Use of a 2-Dose Schedule for Human Papillomavirus Vaccination - Updated Recommendations of the Advisory Committee on Immunization Practices. *MMWR* **65**, 1405-8 (2016).
- ⁷ Stokley, S. et al. Human papillomavirus vaccination coverage among adolescents, 2007-2013, and postlicensure vaccine safety monitoring, 2006-2014--United States. *MMWR Morb Mortal Wkly Rep* **63**, 620-624 (2014).
- ⁸ Naud, P. S. et al. Sustained efficacy, immunogenicity, and safety of the HPV-16/18 AS04-adjuvanted vaccine: final analysis of a long-term follow-up study up to 9.4 years post-vaccination. *Hum Vaccin Immunother* **10**, 2147-2162 (2014).
- ⁹ Joura, E. A. et al. A 9-valent HPV vaccine against infection and intraepithelial neoplasia in women. *N Engl J Med* **372**, 711-723 (2015).
- ¹⁰ Scheller, N. M. et al. Quadrivalent HPV vaccination and risk of multiple sclerosis and other demyelinating diseases of the central nervous system. *JAMA* **313**, 54-61 (2015).
- ¹¹ Scheller, N. M., Pasternak, B., Svanstrom, H. & Hviid, A. Quadrivalent human papillomavirus vaccine and the risk of venous thromboembolism. *JAMA* **312**, 187-188 (2014).
- ¹² Herrero, R. et al. Reduced prevalence of oral human papillomavirus (HPV) 4 years after bivalent HPV vaccination in a randomized clinical trial in Costa Rica. *PLoS One* **8**, e68329 (2013).
- ¹³ Chaturvedi, A.K., et al., *Effect of Prophylactic Human Papillomavirus (HPV) Vaccination on Oral HPV Infections Among Young Adults in the United States*. *J Clin Oncol*, 2018. **36**(3): p. 262-267
- ¹⁴ Lowy DR, Herrero R, Hildesheim A. Primary endpoints for future prophylactic human papillomavirus vaccine trials: towards infection and immunobridging. Participants of the IARC/NCI Workshop on Primary Endpoints for Prophylactic HPV Vaccine Trials. *Lancet Oncol* **16**: e226-33 (2015).
- ¹⁵ Elam-Evans, L. D. et al. National, regional, state, and selected local area vaccination coverage among adolescents aged 13-17 years--United States, 2013. *MMWR Morb Mortal Wkly Rep* **63**, 625-633 (2014).
- ¹⁶ Lu, P. J. et al. Human papillomavirus vaccine initiation and awareness: U.S. young men in the 2010 National Health Interview Survey. *Am J Prev Med* **44**, 330-338 (2013).
- ¹⁷ Jemal, A. et al. Annual Report to the Nation on the Status of Cancer, 1975-2009, featuring the burden and trends in human papillomavirus(HPV)-associated cancers and HPV vaccination coverage levels. *J Natl Cancer Inst* **105**, 175-201 (2013).
- ¹⁸ Tan TQ, Gerbie MV. Perception, Awareness, and Acceptance of Human Papillomavirus Disease and Vaccine Among Parents of Boys Aged 9 to 18 Years. *Clin Pediatr (Phila)* **56**, 737-743 (2017).
- ¹⁹ Finney Rutten LJ, St Sauver JL, Beebe TJ, Wilson PM, Jacobson DJ, Fan C, Breitkopf CR, Vadaparampil ST, MacLaughlin KL, Jacobson RM. Association of both consistency and strength of self-reported clinician recommendation for HPV vaccination and HPV vaccine uptake among 11- to 12-year-old children. *Vaccine* **35**, 6122-6128 (2017).
- ²⁰ O'Rorke, M. A. et al. Human papillomavirus related head and neck cancer survival: a systematic review and meta-analysis. *Oral Oncol* **48**, 1191-1201 (2012).